

# Sago Artikel

*by* Budhi Setiawan

---

**Submission date:** 06-Aug-2022 10:08AM (UTC+0700)

**Submission ID:** 1879339508

**File name:** SAGO\_DRAFT\_MAUSCRIPT\_FINAL\_TEMPLATE.pdf (434.56K)

**Word count:** 6542

**Character count:** 36620

## Review Article

**Physiological response to sago-based foods consumption: a narrative review****Budhi Setiawan<sup>1\*</sup>, Fetriyuna Fetriyuna<sup>2,3</sup>, Sri Murniani Angelina Letsoin<sup>5,6</sup>, Ratna Chrismiari Purwestri<sup>3,7</sup>, Ignasius Radix A.P. Jati<sup>8</sup>**Department of Pharmacology, Faculty of Medicine, Universitas Wijaya Kusuma Surabaya, Dukuh Kupang XXV No. 54, Surabaya 60225, Indonesia<sup>1</sup><sup>2</sup> Department of Food Technology, Faculty of Agro-Industrial Technology, Universitas Padjadjaran, Jln. Raya Bandung-Sumedang Km. 21 Jatinangor 45363, Indonesia<sup>2</sup>Institute of Nutritional Science (140a), University of Hohenheim, Garbenstrasse 30, 70599 Stuttgart, Germany<sup>3</sup><sup>2</sup> Department of Mechanical Engineering, Faculty of Engineering, Czech University of Life Sciences Prague<sup>4</sup>Kamýcká 129, 16500 Praha-Suchdol, Czech Republic<sup>5</sup><sup>2</sup> Faculty of Engineering, University of Musamus, Merauke Regency, 99611, Indonesia<sup>6</sup>  
Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 16500 Praha-Suchdol, Czech Republic<sup>7</sup>Department of Food Technology, Widya Mandala Surabaya Catholic University, Jl. Dinoyo 42-44, Surabaya, 60265, Indonesia<sup>8</sup>

Correspondence author email: budhisetiawan@uwks.ac.id

Phone: +6287751644522

**Abstract**

Sago is a carbohydrate-rich food that contains resistant starch. In some parts of Indonesia, Malaysia, and Papua New Guinea, sago is a common staple food. Studies have been performed to elucidate the physicochemical and structural properties of sago starches. The paper's objective is to review potential physiological responses to sago-based product consumption from previous records. This study is a narrative review of preceding published articles related to sago intake in human subjects research. The literature search was performed through databases with assigned keywords combination and then selection of the articles based on the criteria. The outcomes of this review concluded that sago had lower glycemic index with immediate intestinal absorption. Sago-based products could provide a suitable energy source for sustaining physical performance and promoting faster recovery after exercise, particularly in a warm climate. The prebiotic properties of sago resistant starch have promoted a beneficial composition of gut bacteria, improving the integrity and functionality of the intestinal barrier. Sago starches intake showed positive influences on metabolic indices such as insulin sensitivity and lipid profile. Due to its advantageous physiological effects, thus sago can be an appropriate alternative as an energy source for active healthy individuals as well as conditions like obesity and diabetes.

**Keywords:** Sago; glycemic index; physiological response; resistant starch; carbohydrate; supplement.

## **Respon fisiologis terhadap konsumsi makanan berbahan dasar sago: sebuah tinjauan naratif**

### **Abstrak**

*Sagu adalah makanan kaya karbohidrat yang mengandung pati resisten. Di beberapa daerah di negara-negara seperti Indonesia, Malaysia, dan Papua Nugini, sago adalah makanan pokok yang umum dikonsumsi. Telah dilakukan penelitian untuk menjelaskan sifat fisikokimia dan struktur pati sago. Tujuan tinjauan ini adalah untuk meninjau potensi respon fisiologis terhadap konsumsi produk berbahan dasar sago dari penelitian dengan subyek manusia yang dilakukan sebelumnya. Penelitian ini berupa tinjauan naratif dari beberapa artikel yang telah dipublikasi sebelumnya. Pencarian literatur dilakukan pada beberapa basis data dengan menggunakan kombinasi kata kunci yang telah ditetapkan dan kemudian seleksi artikel didasarkan pada kriteria. Hasil tinjauan ini menyimpulkan bahwa sago memiliki indeks glikemik yang lebih rendah dengan penyerapan di usus yang cepat. Produk berbasis sago menyediakan sumber energi yang dapat mempertahankan kinerja fisik dan mempercepat pemulihan setelah berolahraga, terutama di tempat dengan iklim yang hangat. Sifat prebiotik dari pati resisten sago dapat memfasilitasi komposisi bakteri usus yang bermanfaat, mempertahankan keutuhan dan fungsi dinding usus. Asupan pati sago menunjukkan pengaruh positif pada indeks indeks metabolisme seperti sensitivitas insulin dan profil lipid. Karena efek fisiologisnya yang menguntungkan ini, sago dapat menjadi sebuah alternatif yang tepat sebagai sumber energi untuk individu sehat yang aktif serta beberapa keadaan seperti misalnya obesitas dan diabetes.*

**Kata Kunci:** Sago; indeks glikemik; respon fisiologis; pati resisten; karbohidrat; suplemen.

Received: \_\_\_\_\_ Revised: \_\_\_\_\_ Accepted: \_\_\_\_\_

### **INTRODUCTION**

The Sago palm (*Metroxylon sago*) grows naturally in tropical forests that can produce sago starch extracted from the pith or medulla of the plant's stem (Karim et al., 2008). In Papua New Guinea and Indonesia, sago starch is a staple food that is traditionally prepared by heating stones or by mixing it with hot water (sago porridge, so-called *papeda*) (Metaragakusuma et al., 2016). The significance of sago in the typical diet and as a staple food in the domestic is crucial since it is a food with a high carbohydrate content that is advised to be consumed as a substitute for staple foods. Sago starch usage can aid in a wider variety of foods (Burlingame & Dernini, 2012).

The development of numerous bakery products over the past few decades, including bread, cake, croissants, cookies, and crackers, has raised a demand for wheat flour (Enghiad et al., 2017). Due to increasing consumer awareness of the harmful effects of consuming wheat gluten on health such as wheat allergy, coeliac disease, and non-coeliac gluten sensitivity (Balakireva & Znyatnin Jr, 2016; Czaja-Bulsa, 2015), it opens up an opportunity for sago utilization. Sago flour and sago starch can be applied as components in bread products, primarily as a partial substitution for wheat flour because of their physicochemical characteristics (Konuma et al., 2012). Sago can be used in bakery goods like cake, bread, and brownies in place of wheat flour because it is gluten-free and a rich source of starch (Kumari, 2019).

Sago starch contains approximately 90 percent carbohydrate per 100 grams, which is about similar to tapioca (89 percent) and sweet potato (86 percent), but significantly higher than white rice (34 percent)


or wheat (59 to 71 percent). Furthermore, Sago starch comprises a small proportion of protein and very low-fat content per 100 grams, which are 0.1 percent and 0.01 percent, respectively (Escarnot et al., 2012; Grace & Henry, 2020; Sonia et al., 2015). In order to increase the protein and micronutrient content of a sago-based cuisine, it was later combined with food sources of both animal and plant protein, such as mung beans, soybeans, mushrooms, and fish as the source of fat and minerals (Azkia et al., 2021; Litaay et al., 2021; Mogra & Midha, 2013; Tjokrokusumo et al., 2019).

In Asian diets, carbohydrates are the main source of energy. However, the quantity and quality of carbohydrates are crucial in the management and prevention of diabetes. Foods with a high glycemic index raise blood sugar and insulin levels, which in turn encourage insulin resistance and type 2 diabetes (T2D) through depleting pancreatic  $\beta$  cells (Mohan et al., 2018). The inquiry of whether high carbohydrate proportion in diets has a beneficial or unfavorable effect on cardiometabolic risk factors has sparked interest, especially for T2D. However, the amount of carbohydrates in high carbohydrate diets has no universal definition among researchers, and studies use different standards. Because the definition of high carbohydrate or low carbohydrate diet can be subjective, the labels "higher carbohydrate diet" or "lower carbohydrate diet" seem to be more accurate (Accurso et al., 2008; Feinman et al., 2015; Kodama et al., 2009; Liebman, 2014; Naude et al., 2014). Conversely, the more common staple foods such as white rice and wheat, have more protein (3.9 percent and 10 to 16 percent, respectively) and fat (2.9 percent and 1.1 to 37 percent, respectively) content (Escarnot et al., 2012; Sonia et al., 2015).

Due to the possibility that carbohydrates may have negative effects on postprandial hyperglycemia and aberrant plasma lipid levels, including an increase in triglyceride levels and a decrease in levels of high-density lipoprotein cholesterol (Hyde et al., 2019; Park et al., 2017), the argument over the benefits and drawbacks of a high carbohydrate diet has grown more heated in recent years. However, current findings indicate also that a high good carbohydrate diet does not worsen lipid profiles, notably triglyceride and high-density lipoprotein cholesterol, or glycemic management (Mohan et al., 2018). When made up of high-fiber, low-glycemic index/low-glycemic load foods rather than refined carbohydrates or high-glycemic index foods, a high carbohydrate diet has similar effects to a low carbohydrate diet (Jung & Choi, 2017; Mohan et al., 2018; Schwingshackl & Hoffmann, 2013). Since sago composition is mainly carbohydrate and it has been popular as a staple food as well as an alternative of various flours, thus, the main objective of this paper is to review possible physiological responses of sago-based food consumption in humans.

## METHODS

In order to comprehend the possible impact of sago-based products on physiological responses from human research, a literature review methodology was applied. The literature search used keyword combinations like "sago" AND "supplementation," "sago" AND "glycemic," and "sago" AND "physiologic." The documents were chosen in accordance with the following inclusion criteria: (1) full-text articles from journals that have been published online; (2) publications in English or Indonesian; (3) the articles' relevance to the study's aims; and (4) human subject research. Using the selected keywords, a literature search was conducted from July to August 2022 on Science Direct, PubMed, and Google Scholar databases, and the paper search was limited from 2002 to 2022. All articles obtained from literature search and meeting the criteria were summarized in a table and then discussed using the outcomes of relevant studies.



**Figure 1.** Literature searching strategy and identification via databases using keywords

**RESULT**

The literature search resulted in 147 papers, where most of the papers from the databases were excluded (**Figure 1.**). Based on the more advanced search (i.e., the combination of keywords and accessibility) and thorough checking of the relevancy of the studies, the research objectives, and methodologies for all selected articles, 7 papers in total were included in the review. Four of them were included concerning sago-based product supplementation in exercise and performance, while three articles were reviewed about glycemic index tests (**Table 1.**).

**Table 1.** Summary of the physiological effects of sago-based foods consumption

No.	Methods	General outcomes	References
1.	Seven healthy volunteers (males and females) contributed in the glycemic index test using post-test design. The glycemic indexes of sago noodle and wheat noodle were 26 and 47 respectively with glucose as the standard. In four different types of sago noodles, the resistance starch composition ranged from 7.55 to 9.45 mg/g and wheat resistance starch content was 2.44 mg/g.	Different types of Sago noodle had 3.09 to 3.87 times higher resistant starch composition than wheat noodle. Wheat noodle showed 1.81 times higher glycemic index in comparison with sago noodle.	(Haliza et. al., 2006)
2.	Five clinically healthy individuals participated in the post-test designed glycemic index test. The Glycemic Bread Equivalent (GBE) of foods serves as a representation of a food's relative glycemic	Sago khichdi had faster absorption and lower area under the curve (AUC) of glycemic and insulin responses compared to white bread.	(Singhania & Senray, 2012)

	potency (RGP). Sago khichdi and potatoes had lower glycemic reactions than bread ( $p < 0.05$ ). Potato had an RGP of 52 while sago khichdi had an RGP of 33 per 100 g. Sago khichdi and potatoes had lower overall AUC insulin responses than bread ( $p < 0.05$ ).		
3.	Sago glycemic index test with post-test-controlled design consisted of two groups with 10 healthy volunteers in each group. The glycemic index of pure sago analog rice, analog sago rice with the addition of 5% and 10% red bean were 40.7, 48.3, and 50.4, respectively.	Pure sago analog rice and analog sago rice with red bean flour have shown a low glycemic index.	(Wahjuningsih et al. 2016)
4.	Prospective experimental post-test design with twelve healthy male subjects. The plasma glucose area under the curve (AUC) of sago gel was higher than white bread. The plasma insulin AUC of sago gel was higher than white bread and sago porridge. Sago gel, sago porridge, and sago paste showed comparable glycemic responses. However, the insulin response of sago porridge was higher than sago paste and sago gel.	Supplementation of Sago paste and sago porridge is useful in pre and during exercise to improve performance. Sago gel intake during recovery can be effective during a subsequent endurance activity.	(Ahmad et al., 2009)
5.	A randomized, double-blind placebo-controlled crossover design study with 8 recreational male cyclists as participants. Times to exhaustion after placebo, sago, and sago-soy consumption were $4.09 \pm 1.28$ , $5.49 \pm 1.20$ , and $7.53 \pm 2.02$ min, respectively. Sago-soy supplementation improved endurance during high-intensity cycling by 84% (44–140%; $p < 0.001$ ) and by 37% (15–63%; $p < 0.05$ ) compared to placebo and sago, respectively.	A combination of sago and soy protein consumption could delay fatigue in high-intensity cycling.	(Ghosh et al., 2010)
6.	Eight trained male cyclists participated in a crossover design. Dur-plasma Sago's glucose increased during exercise more than the control's did ( $p < 0.05$ ). Substrate oxidation rates, however, remained constant ( $p > 0.05$ ). Higher plasma sodium levels ( $2 \pm 2$ mmol/liter) were observed, together with lower whole-body sweat loss ( $544 \pm 636$ g) and decreased plasma volume contraction (all $p < 0.05$ ). Dur-Sago caused an increase in heart rate ( $5 \pm 13$ beats/min), but performance was comparable to control ( $+0.9 \pm 2.3\%$ , $p > 0.05$ ).	Sago feeding during exercise with heat stress resulted in some "beneficial" physiological reactions, but post-prandial performance was unaffected.	(Jusoh et al., 2016a)
7.	Crossover design and the subject included 8 trained male cyclists. Average power output was not different between trials during the	Sago consumption during recovery from exercise in a warm, humid environment enhances recovery of	(Jusoh et al., 2016b)

---

<p>first exercise (286 ± 67 vs. 281 ± 59 W), however in the second exercise, the sago group was higher than the control (283 ± 60 vs. 274 ± 61 W, p&lt; 0.05)</p>	<p>performance in following exercise when compared to a water-only control.</p>
---	---

---

## DISCUSSION

In 1981, the glycemic index (GI) was introduced for the first time (Jenkins et al., 1981). Groups of 5 to 10 healthy fasting participants were given particular food individually to see how it affected their blood glucose levels. Blood glucose levels were assessed over a period of two hours and expressed as a proportion of the area under the glucose response curve for a given amount of the carbohydrate (Jenkins et al., 1981). It refers to the degree and duration of blood glucose elevation that occurs after a 12-hour fast in response to a particular carbohydrate intake ratio to a standard carbohydrate, typically glucose or white bread. Glycemic index is divided into three categories: Low GI (below 56), Medium GI (between 56 and 69), and High GI (beyond 69) on a scale from 0 to 100 (Augustin et al., 2015). It has been shown that sago had a lower glycemic index value (40.7, low level) (Wahjuningsih et al., 2016) compared to the other staple foods, for instance, white rice (61–87, medium to high level) and sweet potato (77, medium) (Osman et al., 2021). Other studies reported the values of sago glycemic index using different settings were 27 (Haliza et al., 2006) and 33 bread equivalent per 100 g (Singhania & Senray, 2012). However, the three commonly used glycemic index categories may be incorrectly assigned to foods due to variations in GI methodologies. It could be related to variations in baseline HbA1c and insulin index, indicating that even in people with normoglycemia, longer-term glycemic control and insulin responsiveness have an impact on the GI score (Matthan et al., 2016).

In human studies, sago has shown preliminary evidence for a beneficial effect on exercise performance (Table 2.). Despite sago having a lower glycemic index, sago-based products provide a higher glycemic response that can improve physical performance during exercise (Ghosh et al., 2010; Matthan et al., 2016; Wahjuningsih et al., 2016). Sago-based meals have demonstrated higher glycemic and insulinemic responses when compared to white bread, and therefore, their intake may improve performance in prolonged exercise (Ahmad et al., 2009). Hence, sago could be an option as a food or supplement for athletes or people in physical exercise to maintain performance and accelerate recovery. It could be due to starch-based foods that are high amylopectin content (e.g., sago and potatoes) speed up the absorption of sugar, supplying the cells' need for energy immediately (Singhania & Senray, 2012).

Sago is more likely to be suitable as a supplement before and during exercise because of its physical and chemical characteristics. For example, sago consumption during exercise showed a beneficial physiologic response (Jusoh et al., 2016a). Also, sago consumption in a recovery time could improve performance during a subsequent exercise period compared to a water-only control (Jusoh et al., 2016b). Nevertheless, further study is required to compare the findings to other known carbohydrate sources. Meanwhile, research was also performed by combining sago with soy protein to improve the protein content of the sago-based product and its functionality. Sago and soy protein consumed after moderately strenuous cycling exercise improved subsequent high-intensity endurance capacity when compared to sago alone and a placebo (Ghosh et al., 2010).

Resistant starch (RS) refers to all types of starch that are digested in the colon to produce short chain fatty acids but are inaccessible to human digestive enzymes (Ashwar et al., 2016). Table 2. describes that RS can be divided into five types (RS1–RS5), some of which are created or altered commercially and added to food as functional ingredients (types 2, 3, and 4), while others occur naturally in foods (types 1, 2, 3, and 5) (Lockyer & Nugent, 2017; Patterson et al., 2020; Tian & Sun, 2020). On the other hand, prebiotics are known as nutrients or food elements that resist digestion in the upper gastrointestinal tract,

fermented by the intestinal microbiota, and on a selective basis, stimulating the growth and/or activity of the beneficial intestinal bacteria. An essential component of the concept is that it must improve host health (Zaman & Sarbini, 2016). Resistant starch type 3 (RS3) can be produced from different parts (top and bottom) of *Metroxylon sago* with different growth stages (Abd Rashid et al., 2020). Produced from *Metroxylon sago* starch, sago resistant starch type III (RS3) has been shown resistant to digestion by stomach acids and enzymes. Additionally, sago RS3 displayed prebiotic properties such as inducing beneficial gut bacteria (e.g., *Bifidobacterium sp.*, *Lactobacillus bulgaricus*, *Clostridium butyricum*, *Eubacterium rectale*) growth with short-chain fatty acid released even on par with those of commercial prebiotics (Purwani et al., 2012; Zi-Ni et al., 2015; Zi-Ni & Rosma, 2020).

**Table 2.** Classification of types of resistant starch (RS), characteristics, and food sources (Ashwar et al., 2016; Lockyer & Nugent, 2017; Raigond et al., 2015)

RS types	Characteristics	Food sources
RS1	Because of the physical defense provided by protein matrices and cell walls, which thwart digesting enzymes.	Whole or partly milled grains, legumes, and seeds.
RS2	Starches are resistant to enzymatic hydrolysis because of their crystalline structure.	Green bananas, raw potatoes, high-amylose maize, and ginkgo starch.
RS3	When starchy foods are cooked and chilled, retrograded starch is created. The establishment of hydrogen bonds during retrogradation allows the polymer chains to reassociate and form double helices.	Potatoes that have been cooked and then allowed to cool, crusty bread, pasta, rice, cornflakes, and food items that have undergone repeated moist heat treatment.
RS4	Modified starch that has undergone chemical alterations, such as esterification, etherification, or cross-linking. Starch cannot be digested due to chemical changes that limit enzyme access and create atypical links.	Foods that have chemically modified starches (e.g., bread and cakes).
RS5	The foods contain amylose-lipid complexes, which can be produced artificially and added to foods or develop naturally during processing and reform after cooking.	Foods with high amylose-lipid complex content.

Besides sago RS3, sago fronds can be enzymatically hydrolyzed to produce sago frond sugar (SFS), which contains 16 to 18 percent cellobiose, known as a type of prebiotic sugar (Dyg Salwani et al., 2018). The sago starch oligosaccharides have been used to enhance the growth of lactic acid bacteria (*Lactobacillus bulgaricus* and *Streptococcus thermophilus*) in homemade yogurt, with the potential to be developed as prebiotic food (Shima et al., 2012). Traditionally processed sago starch (*Lemantak*) had resistant starch content at 62.61 percent and could be used as fermentation substrate for *Lactobacillus acidophilus* and *Bifidobacterium animalis* at 37°C for 24 hours in anaerobic conditions (Arshad et al., 2018). Hence, these abovementioned in vitro study outcomes confirmed the prebiotic properties of sago.

Resistant starch is well known for its role in regulating obesity and T2D through the modification of gut microbiota composition. Starch-degrading enzymes and intestinal metabolites are produced as a result of the change of a specific microbial composition, which enhances the function of the healthy gut barrier (Liu et al., 2020). Gut microbial fermentation of resistant starch in the large bowel increases the short-chain fatty acids (SCFA) composition such as acetate, propionate, and butyrate (Liu et al., 2020; Topping & Clifton, 2001). By controlling luminal pH and mucus production, short-chain fatty acids are crucial for maintaining the integrity of the gut. Additionally, they fuel epithelial cells and control mucosal immune response (Blaak et al., 2020). It has been shown that resistant starch might reduce hyperglycemia, hyperinsulinemia, and hyperlipidemia. This can be due to restricting gluconeogenesis, increasing



glycogenesis, maintaining glucose and lipid homeostasis, and ameliorating pancreatic dysfunction (Meenu & Xu, 2019). In addition, studies have shown that resistant starch intervention may improve the serum level of inflammatory biomarkers such as interleukin-6 and tumor necrosis factor-alpha (Vahdat et al., 2020), thus it is suggested that the resistant starch can reduce the inflammatory incidence.

In animal model studies, sago-based products intervention has demonstrated potential beneficial effects on metabolic indices. Sago resistant starch has shown potential as a strategy for obesity and overweight control in obese rats. Sago resistant starch promoted satiety, lowered visceral adipose tissue, and improved hepatic lipid content (Thompson et al., 2021). Anti-obesity effect could be due to prebiotic properties of sago starch which had resulted in higher *Bifidobacterium spp.* and *Lactobacillus spp.* growth, significant increase of total concentration fecal and caecum short chain fatty acids and lower fat accumulation in liver (Laang, 2018). In line with aforementioned outcomes, sago consumption has played significant role in diabetic animal models. For examples, sago-based products intervention has improved insulin sensitivity and lipid profile (Wahjuningsih et al., 2018) as well as stimulated better pancreatic Langerhans  $\beta$ - cell function. The mechanisms involved are absorption reduction, insulin expression of pancreatic  $\beta$ -cell elevation, and total amount of short chain fatty acids rise (Wahjuningsih et al., 2020). Furthermore, modified sago resistant starches (RS2 and RS3 types) intake could down/upregulate genes which are related to regulation and enzymatic pathway of glucose and glycogen metabolism in liver that have important role in diabetes management (Lokman et al., 2021). Additionally, besides metabolic benefits, sago supplementation has been shown advantageous effects on antioxidant activity indices such as liver superoxide dismutase, catalase, liver  $\alpha$ -tocopherol (Hirao & Igarashi, 2003).

## CONCLUSION

This review revealed possible influences of sago starch consumption on human health, that can be further study direction. Sago resistant starch provides favorable physiological responses that can be due to its lower glycemic index and faster absorption. Furthermore, prebiotic properties of Sago resistant starch induce beneficial intestinal microbiota composition that result in increased SFCA levels and better intestinal epithelium protection. Consumption of sago starches has a favorable impact on metabolic parameters like improved pancreatic  $\beta$ -cells and insulin functions as well as lipid panel. Sago-based foods make a good source of supplements for sustaining physical performance and hastening recovery in period of peri-exercise. The physical and chemical properties of sago, which is frequently grown and consumed in warm, humid climates, make it suitable as a supplement for physical work out. Sago can be a suitable alternative as an energy source for active healthy people as well as disorders like obesity and diabetes because of its beneficial physiological benefits.

## ACKNOWLEDGEMENT

The authors would like to thank the Czech University of Life Sciences Prague for financing the project "Advanced research supporting the forestry and wood-processing sector's adaptation to global change and the fourth industrial revolution" (EVA 4.0). The chance to perform this review study was made possible by the support of the Directorate of Research and Community Services at the Universitas Padjadjaran, which is also acknowledged by the authors. Regina Magaa Vázquez and Monika Neuss provided assistance with the Humboldt reloaded project. Dhea Ayu Putri assisted with the literature review assignment. Yus Widarko from the University of Musamus shared information about his research in Tambat, Merauke.

**REFERENCES**

- Abd Rashid, R. S., Dos Mohamed, A. M., Achudan, S. N., & Mittis, P. (2020). Physicochemical properties of resistant starch type III from sago starch at different palm stages. *Materials Today: Proceedings*, 31, 150–154.
- Accurso, A., Bernstein, R. K., Dahlqvist, A., Draznin, B., Feinman, R. D., Fine, E. J., Gleed, A., Jacobs, D. B., Larson, G., & Lustig, R. H. (2008). Dietary carbohydrate restriction in type 2 diabetes mellitus and metabolic syndrome: Time for a critical appraisal. *Nutrition & Metabolism*, 5(1), 1–8. <https://doi.org/10.1186/1743-7075-5-9>
- Ahmad, H., Singh, R., & Ghosh, A. K. (2009). Glycaemic & insulinaemic responses in men at rest following sago meal. *The Indian Journal of Medical Research*, 130(2), 160–165.
- Arshad, N. H., Zaman, S. A., Rawi, M. H., & Sarbini, S. R. (2018). Resistant starch evaluation and in vitro fermentation of *lemantak* (native sago starch), for prebiotic assessment. *International Food Research Journal*, 25(3).
- Ashwar, B. A., Gani, A., Shah, A., Wani, I. A., & Masoodi, F. A. (2016). Preparation, health benefits and applications of resistant starch—A review. *Starch-Stärke*, 68(3–4), 287–301. <https://doi.org/10.1002/star.201500064>
- Augustin, L. S., Kendall, C. W., Jenkins, D. J., Willett, W. C., Astrup, A., Barclay, A. W., Björck, I., Brand-Miller, J. C., Brighenti, F., & Buyken, A. E. (2015). Glycemic index, glycemic load and glycemic response: An International Scientific Consensus Summit from the International Carbohydrate Quality Consortium (ICQC). *Nutrition, Metabolism and Cardiovascular Diseases*, 25(9), 795–815. <https://doi.org/10.1016/j.numecd.2015.05.005>
- Azka, M. N., Wahjuningsih, S. B., & Wibowo, C. H. (2021). The nutritional and functional properties of noodles prepared from sorghum, mung bean and sago flours. *Food Research*, 5(S2), 65–69. [https://doi.org/10.26656/fr.2017.5\(S2\).002](https://doi.org/10.26656/fr.2017.5(S2).002)
- Balakireva, A. V., & Zamyatnin Jr, A. A. (2016). Properties of gluten intolerance: Gluten structure, evolution, pathogenicity and detoxification capabilities. *Nutrients*, 8(10), 644. <https://doi.org/10.3390/nu8100644>
- Blaak, E. e., Canfora, E. e., Theis, S., Frost, G., Groen, A. k., Mithieux, G., Nauta, A., Scott, K., Stahl, B., van Harsselaar, J., van Tol, R., Vaughan, E. e., & Verbeke, K. (2020). Short chain fatty acids in human gut and metabolic health. *Beneficial Microbes*, 11(5), 411–455. <https://doi.org/10.3920/BM2020.0057>
- Burlingame, B., & Dernini, S. (2012). Biodiversity and sustainable diets united against hunger 3–5 November 2010 FAO Headquarters, Rome.
- Czaja-Bulsa, G. (2015). Non coeliac gluten sensitivity—A new disease with gluten intolerance. *Clinical Nutrition*, 34(2), 189–194.
- Dyg Salwani, A. A., Ahmad, M. N., & Bujang, K. (2018). Maximising production of prebiotic sugar (cellobiose) from sago frond. *Malaysian Applied Biology Journal*, 47(1), 89–95.
- Enghiad, A., Ufer, D., Countryman, A. M., & Thilmany, D. D. (2017). An overview of global wheat market fundamentals in an era of climate concerns. *International Journal of Agronomy*, 2017.
- Escarnot, E., Jacquemin, J.-M., Agneessens, R., & Paquot, M. (2012). Comparative study of the content and profiles of macronutrients in spelt and wheat, a review. *BASE*. <https://popups.uliege.be/1780-4507/index.php?id=8700>

- Feinman, R. D., Pogozelski, W. K., Astrup, A., Bernstein, R. K., Fine, E. J., Westman, E. C., Accurso, A., Frassetto, L., Gower, B. A., & McFarlane, S. I. (2015). Dietary carbohydrate restriction as the first approach in diabetes management: Critical review and evidence base. *Nutrition*, 31(1), 1–13. <https://doi.org/10.1016/j.nut.2014.06.011>
- Ghosh, A. K., Rahaman, A. A., & Singh, R. (2010). Combination of sago and soy-protein supplementation during endurance cycling exercise and subsequent high-intensity endurance capacity. *International Journal of Sport Nutrition and Exercise Metabolism*, 20(3), 216–223. <https://doi.org/10.1123/ijsnem.20.3.216>
- Grace, N. C., & Henry, C. J. (2020). The physicochemical characterization of unconventional starches and flours used in Asia. *Foods*, 9(2), 182. <https://doi.org/10.3390/foods9020182>
- Haliza, W., Purwani, E. Y., & Yuliani, S. (2006). Evaluation Of Enzymatically Resistant Starch and Glycemix Index of Sago Noodle. *Jurnal Teknologi dan Industri Pangan*, 17(2), 150–150.
- Hirao, K., & Igarashi, K. (2003). Effects of Sago Starch Content in the Diet on Lipid Peroxidation and Antioxidative Enzyme Activities in Rats. *Journal of Nutritional Science and Vitaminology*, 49(1), 76–79. <https://doi.org/10.3177/jnsv.49.76>
- Hyde, P. N., Sapper, T. N., Crabtree, C. D., LaFountain, R. A., Bowling, M. L., Buga, A., Fell, B., McSwiney, F. T., Dickerson, R. M., Miller, V. J., Scandling, D., Simonetti, O. P., Phinney, S. D., Kraemer, W. J., King, S. A., Krauss, R. M., & Volek, J. S. (2019). Dietary carbohydrate restriction improves metabolic syndrome independent of weight loss. *JCI Insight*, 4(12), 128308. <https://doi.org/10.1172/jci.insight.128308>
- Jenkins, D. J., Wolever, T. M., Taylor, R. H., Barker, H., Fielden, H., Baldwin, J. M., Bowling, A. C., Newman, H. C., Jenkins, A. L., & Goff, D. V. (1981). Glycemic index of foods: A physiological basis for carbohydrate exchange. *The American Journal of Clinical Nutrition*, 34(3), 362–366. <https://doi.org/10.1093/ajcn/34.3.362>
- Jung, C.-H., & Choi, K. M. (2017). Impact of high-carbohydrate diet on metabolic parameters in patients with type 2 diabetes. *Nutrients*, 9(4), 322. <https://doi.org/10.3390/nu9040322>
- Jusoh, M. R. C., Stannard, S. R., & Mündel, T. (2016a). Physiologic and performance effects of sago supplementation before and during cycling in a warm-humid environment. *Temperature*, 3(2), 318–327. <https://doi.org/10.1080%2F23328940.2016.1159772>
- Jusoh, M. R. C., Stannard, S. R., & Mündel, T. (2016b). Sago supplementation for recovery from cycling in a warm-humid environment and its influence on subsequent cycling physiology and performance. *Temperature*, 3(3), 444–454. <https://doi.org/10.1080%2F23328940.2016.1179382>
- Karim, A. A., Tie, A. P.-L., Manan, D. M. A., & Zaidul, I. S. M. (2008). Starch from the sago (*Metroxylon sagu*) palm tree—Properties, prospects, and challenges as a new industrial source for food and other uses. *Comprehensive Reviews in Food Science and Food Safety*, 7(3), 215–228. <https://doi.org/10.1111/j.1541-4337.2008.00042.x>
- Kodama, S., Saito, K., Tanaka, S., Maki, M., Yachi, Y., Sato, M., Sugawara, A., Totsuka, K., Shimano, H., & Ohashi, Y. (2009). Influence of fat and carbohydrate proportions on the metabolic profile in patients with type 2 diabetes: A meta-analysis. *Diabetes Care*, 32(5), 959–965. <https://doi.org/10.2337/dc08-1716>
- Konuma, H., Rolle, R., & Boromthanasarat, S. (2012). Adding value to underutilized food resources: Substituting wheat flour with sago starch in cookie formulations. *Journal of Agricultural Technology*, 8(3), 1077.

- Laang, M. S. T. (2018). Potential of native sago starch as prebiotic with an anti-obesity functionality. Potential of native sago starch as prebiotic with an anti-obesity functionality. *Master thesis*, Universiti Putra Malaysia.
- Liebman, M. (2014). When and why carbohydrate restriction can be a viable option. *Nutrition*, 30(7–8), 748–754. <https://doi.org/10.1016/j.nut.2013.11.021>
- Litaay, C., Indriati, A., & Mayasti, N. K. I. (2021). Fortification of sago noodles with fish meal skipjack tuna (*Katsuwonus pelamis*). *Food Science and Technology*. <https://doi.org/10.1590/fst.46720>
- Liu, H., Zhang, M., Ma, Q., Tian, B., Nie, C., Chen, Z., & Li, J. (2020). Health beneficial effects of resistant starch on diabetes and obesity via regulation of gut microbiota: A review. *Food & Function*, 11(7), 5749–5767. <https://doi.org/10.1039/D0FO00855A>
- Lockyer, S., & Nugent, A. P. (2017). Health effects of resistant starch. *Nutrition Bulletin*, 42(1), 10–41.
- Lokman, E. F., Aziz, S. M. A., Ibrahim, A. S., Yunus, N., Seruji, A. Z. R. A., & Bugam, S. H. (2021). Hepatic Glucose Regulations by Sago (*Metroxylon sago*) Resistant Starch in Diabetic Goto Kakizaki Rat. *International Journal of Biomedical Science*, 17(3), 28–33.
- Matthan, N. R., Ausman, L. M., Meng, H., Tighiouart, H., & Lichtenstein, A. H. (2016). Estimating the reliability of glycemic index values and potential sources of methodological and biological variability. *The American Journal of Clinical Nutrition*, 104(4), 1004–1013. <https://doi.org/10.3945/ajcn.116.137208>
- Meenu, M., & Xu, B. (2019). A critical review on anti-diabetic and anti-obesity effects of dietary resistant starch. *Critical Reviews in Food Science and Nutrition*, 59(18), 3019–3031. <https://doi.org/10.1080/10408398.2018.1481360>
- Metaragakusuma, A. P., Katsuya, O., & Bai, H. (2016). An Overview of the Traditional use of sago for Sago-based Food Industry in Indonesia. *KnE Life Sciences*, 119–124. <https://dx.doi.org/10.18502/cls.v3i3.382>
- Mogra, R., & Midha, S. (2013). Value addition of traditional wheat flour vermicelli. *Journal of Food Science and Technology*, 50(4), 815–820. <https://doi.org/10.1007/s13197-011-0403-3>
- Mohan, V., Unnikrishnan, R., Shobana, S., Malavika, M., Anjana, R. M., & Sudha, V. (2018). Are excess carbohydrates the main link to diabetes & its complications in Asians? *The Indian Journal of Medical Research*, 148(5), 531. [https://doi.org/10.4103%2Fijmr.IJMR\\_1698\\_18](https://doi.org/10.4103%2Fijmr.IJMR_1698_18)
- Naude, C. E., Schoonees, A., Senekal, M., Young, T., Garner, P., & Volmink, J. (2014). Low carbohydrate versus isoenergetic balanced diets for reducing weight and cardiovascular risk: A systematic review and meta-analysis. *PloS One*, 9(7), e100652. <https://doi.org/10.1371/journal.pone.0100652>
- Osman, M. H., Yusof, B. N., & Ismail, A. (2021). Glycaemic index and glycaemic load of foods and food products in Malaysia: A review. *International Food Research Journal*, 28(2), 217–229.
- Park, S., Ahn, J., Kim, N.-S., & Lee, B.-K. (2017). High carbohydrate diets are positively associated with the risk of metabolic syndrome irrespective to fatty acid composition in women: The KNHANES 2007–2014. *International Journal of Food Sciences and Nutrition*, 68(4), 479–487. <https://doi.org/10.1080/09637486.2016.1252318>
- Patterson, M. A., Maiya, M., & Stewart, M. L. (2020). Resistant starch content in foods commonly consumed in the United States: A narrative review. *Journal of the Academy of Nutrition and Dietetics*, 120(2), 230–244. <https://doi.org/10.1016/j.jand.2019.10.019>
- Purwani, E. Y., Purwadaria, T., & Suhartono, M. T. (2012). Fermentation RS3 derived from sago and rice starch with *Clostridium butyricum* BCC B2571 or *Eubacterium rectale* DSM 17629. *Anaerobe*, 18(1), 55–61. <https://doi.org/10.1016/j.anaerobe.2011.09.007>

- Raigond, P., Ezekiel, R., & Raigond, B. (2015). Resistant starch in food: A review. *Journal of the Science of Food and Agriculture*, 95(10), 1968–1978.
- Schwingshackl, L., & Hoffmann, G. (2013). Long-term effects of low glycemic index/load vs. High glycemic index/load diets on parameters of obesity and obesity-associated risks: A systematic review and meta-analysis. *Nutrition, Metabolism and Cardiovascular Diseases*, 23(8), 699–706. <https://doi.org/10.1016/j.numecd.2013.04.008>
- Shima, A. R., Salina, H. F., Masniza, M., & Atiqah, A. H. (2012). Viability of Lactic Acid Bacteria in Home Made Yogurt Containing Sago Starch Oligosaccharides. 12(01), 58-62.
- Singhania, P. R., & Senray, K. (2012). Glycemic response to amylopectin rich starch present in common fasting foods of India. *Nutrition & Food Science*, 42(3), 196–203. <https://doi.org/10.1108/00346651211228496>
- Sonia, S., Witjaksono, F., & Ridwan, R. (2015). Effect of cooling of cooked white rice on resistant starch content and glycemic response. *Asia Pacific Journal of Clinical Nutrition*, 24(4), 620–625.
- Thompson, M. S., Dahari, S. I., Shamsuddin, M. S., Rashed, A., & Sarbini, S. R. (2021). Effects of sago starch on body weight, food intake, caecum short chain fatty acids, adipose tissue, and hepatic lipid content of fat-induced Sprague Dawley rats. *International Food Research Journal*, 28(5), 1057–1066. <https://doi.org/10.47836/ifrj.28.5.19>
- Tian, S., & Sun, Y. (2020). Influencing factor of resistant starch formation and application in cereal products: A review. *International Journal of Biological Macromolecules*, 149, 424–431. <https://doi.org/10.1016/j.ijbiomac.2020.01.264>
- Tjokrokusumo, D., Octaviani, F. C., & Saragih, R. (2019). Fortification of Mung bean (*Vigna radiata*) and Ear mushroom (*Auricularia auricula-judae*) in dried sago noodles. *Journal of Microbial Systematics and Biotechnology*, 1(2), 34–40. <https://doi.org/10.37604/jmsb.v1i2.30>
- Topping, D. L., & Clifton, P. M. (2001). Short-chain fatty acids and human colonic function: Roles of resistant starch and nonstarch polysaccharides. *Physiological Reviews*, 81(3), 1031–1064. <https://doi.org/10.1152/physrev.2001.81.3.1031>
- Vahdat, M., Hosseini, S. A., Khalatbari Mohseni, G., Heshmati, J., & Rahimlou, M. (2020). Effects of resistant starch interventions on circulating inflammatory biomarkers: A systematic review and meta-analysis of randomized controlled trials. *Nutrition Journal*, 19, 33. <https://doi.org/10.1186/s12937-020-00548-6>
- Wahjuningsih, S. B., Haslina, H., & Marsono, M. (2018). Hypolipidaemic Effects of High Resistant Starch Sago and Red Bean Flour- based Analog Rice on Diabetic Rats. *Materia Socio-Medica*, 30(4), 232–239. <https://doi.org/10.5455/msm.2018.30.232-239>
- Wahjuningsih, S. B., Haslina, H., & Putranto, A. T. (2020). Effect of Sago Analogue Rice and Red Bean Diet to the Improvement of  $\beta$ -cell Pancreas in Streptozotocin-Nicotinamide (STZ-NA) Induced Diabetic Rats. *Current Research in Nutrition and Food Science*, 8(2), 667. <http://dx.doi.org/10.12944/CRNFSJ.8.2.32>
- Wahjuningsih, S. B., Marsono, Y., Praseptiangga, D., & Haryanto, B. (2016). Resistant Starch Content and Glycaemic Index of Sago (*Metroxylon spp.*) Starch and Red Bean (*Phaseolus vulgaris*) Based Analogue Ric. *Science Alert*, 15(7), 667–672. <https://doi.org/10.3923/pjn.2016.667.672>
- Zaman, S. A., & Sarbini, S. R. (2016). The potential of resistant starch as a prebiotic. *Critical Reviews in Biotechnology*, 36(3), 578–584. <https://doi.org/10.3109/07388551.2014.993590>
- Zi-Ni, T., & Rosma, A. (2020). Selective Fermentation and Prebiotic Index of Sago (*Metroxylon sago*) Resistant Starches Type III by Rat Fecal Cultures. IOP Conference Series: Materials Science and Engineering, 716, 012002. <https://doi.org/10.1088/1757-899X/716/1/012002>

Zi-Ni, T., Rosma, A., Napisah, H., Karim, A. A., & Liong, M.-T. (2015). Characteristics of *Metroxylon sagu* resistant starch type III as prebiotic substance. *Journal of Food Science*, 80(4), H875–H882. <https://doi.org/10.1111/1750-3841.12817>

# Sago Artikel

---

## ORIGINALITY REPORT

---

10%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

1%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://journal.uwks.ac.id">journal.uwks.ac.id</a> Internet Source	3%
2	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	2%
3	<a href="http://lib.bioinfo.pl">lib.bioinfo.pl</a> Internet Source	1%
4	"Sago Palm", Springer Science and Business Media LLC, 2018 Publication	1%
5	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1%
6	Mohd Rahimi Che Jusoh, Stephen R. Stannard, Toby Mündel. "Sago supplementation for recovery from cycling in a warm-humid environment and its influence on subsequent cycling physiology and performance", Temperature, 2016 Publication	1%
7	Polysaccharides, 2015. Publication	1%

---

8

Renbing Qin, Jinglin Yu, Yufang Li, Les Copeland, Shuo Wang, Shujun Wang.  
"Structural Changes of Starch–Lipid Complexes during Postprocessing and Their Effect on In Vitro Enzymatic Digestibility",  
Journal of Agricultural and Food Chemistry,  
2019

Publication

1 %

9

[onlinelibrary.wiley.com](https://onlinelibrary.wiley.com)

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On