

LAMPIRAN

Lampiran 1. Hasil Analisis Univariat

Usia

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Anak-anak	16	32.0	32.0	32.0
	Dewasa	31	62.0	62.0	94.0
	Lansia	3	6.0	6.0	100.0
	Total	50	100.0	100.0	

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	27	54.0	54.0	54.0
	Perempuan	23	46.0	46.0	100.0
	Total	50	100.0	100.0	

IMT

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Underweight	13	26.0	26.0	26.0
	Normal	21	42.0	42.0	68.0
	Overweight	11	22.0	22.0	90.0
	Obese	5	10.0	10.0	100.0
	Total	50	100.0	100.0	

Kondisi Imunokompromais

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	26	52.0	52.0	52.0
	Tidak	24	48.0	48.0	100.0
	Total	50	100.0	100.0	

Riwayat TB Paru

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	29	58.0	58.0	58.0
	Tidak	21	42.0	42.0	100.0
	Total	50	100.0	100.0	

Lokasi Lesi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Cervical	2	4.0	4.0	4.0
	Thoracal	17	34.0	34.0	38.0
	Lumbal	21	42.0	42.0	80.0
	Multiple	10	20.0	20.0	100.0
	Total	50	100.0	100.0	

Tingkat Keparahan Spondilitis TB

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ringan	13	26.0	26.0	26.0
	Sedang	16	32.0	32.0	58.0
	Berat	21	42.0	42.0	100.0
	Total	50	100.0	100.0	

Tata Laksana

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Konservatif	8	16.0	16.0	16.0
	Operatif	42	84.0	84.0	100.0
	Total	50	100.0	100.0	

Lampiran 2. Hasil Analisis Bivariat

Usia * Tingkat Keparahan Spondilitis TB Crosstabulation

			Tingkat Keparahan Spondilitis TB			
			Ringan	Sedang	Berat	Total
Usia	Anak-anak	Count	3	7	6	16
		Expected Count	4.2	5.1	6.7	16.0
		% within Usia	18.8%	43.8%	37.5%	100.0%
	Dewasa	Count	9	8	14	31
		Expected Count	8.1	9.9	13.0	31.0
		% within Usia	29.0%	25.8%	45.2%	100.0%
	Lansia	Count	1	1	1	3
		Expected Count	.8	1.0	1.3	3.0
		% within Usia	33.3%	33.3%	33.3%	100.0%
Total	Count	13	16	21	50	
	Expected Count	13.0	16.0	21.0	50.0	
	% within Usia	26.0%	32.0%	42.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	1.763 ^a	4	.779
Likelihood Ratio	1.749	4	.782
Linear-by-Linear Association	.081	1	.776
N of Valid Cases	50		

a. 4 cells (44.4%) have expected count less than 5. The minimum expected count is .78.

Usia * Tingkat Keparahan Spondilitis TB Crosstabulation

			Tingkat Keparahan Spondilitis TB			
			Ringan	Sedang	Berat	Total
Usia	Risiko Tinggi	Count	4	8	7	19
		Expected Count	4.9	6.1	8.0	19.0
		% within Usia	21.1%	42.1%	36.8%	100.0%
	Risiko Rendah	Count	9	8	14	31
		Expected Count	8.1	9.9	13.0	31.0
		% within Usia	29.0%	25.8%	45.2%	100.0%
Total	Count	13	16	21	50	

Expected Count	13.0	16.0	21.0	50.0
% within Usia	26.0%	32.0%	42.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	1.461 ^a	2	.482
Likelihood Ratio	1.444	2	.486
Linear-by-Linear Association	.000	1	.989
N of Valid Cases	50		

a. 1 cells (16.7%) have expected count less than 5. The minimum expected count is 4.94.

Jenis Kelamin * Tingkat Keparahan Spondilitis TB Crosstabulation

		Tingkat Keparahan Spondilitis TB			Total	
		Ringan	Sedang	Berat		
Jenis Kelamin	Laki-laki	Count	6	11	10	27
		Expected Count	7.0	8.6	11.3	27.0
		% within Jenis Kelamin	22.2%	40.7%	37.0%	100.0%
	Perempuan	Count	7	5	11	23
		Expected Count	6.0	7.4	9.7	23.0
		% within Jenis Kelamin	30.4%	21.7%	47.8%	100.0%
Total	Count	13	16	21	50	
	Expected Count	13.0	16.0	21.0	50.0	
	% within Jenis Kelamin	26.0%	32.0%	42.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	2.068 ^a	2	.356
Likelihood Ratio	2.110	2	.348
Linear-by-Linear Association	.012	1	.912
N of Valid Cases	50		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 5.98.

IMT * Tingkat Keparahan Spondilitis TB Crosstabulation

		Tingkat Keparahan Spondilitis TB			Total	
		Ringan	Sedang	Berat		
IMT	Underweight	Count	0	6	7	13
		Expected Count	3.4	4.2	5.5	13.0
		% within IMT	0.0%	46.2%	53.8%	100.0%
	Normal	Count	9	7	5	21
		Expected Count	5.5	6.7	8.8	21.0
		% within IMT	42.9%	33.3%	23.8%	100.0%
	Overweight	Count	0	2	9	11
		Expected Count	2.9	3.5	4.6	11.0
		% within IMT	0.0%	18.2%	81.8%	100.0%
Obese	Count	4	1	0	5	
	Expected Count	1.3	1.6	2.1	5.0	
	% within IMT	80.0%	20.0%	0.0%	100.0%	
Total	Count	13	16	21	50	
	Expected Count	13.0	16.0	21.0	50.0	
	% within IMT	26.0%	32.0%	42.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	24.191 ^a	6	.000
Likelihood Ratio	29.558	6	.000
Linear-by-Linear Association	1.858	1	.173
N of Valid Cases	50		

a. 8 cells (66.7%) have expected count less than 5. The minimum expected count is 1.30.

IMT * Tingkat Keparahan Spondilitis TB Crosstabulation

		Tingkat Keparahan Spondilitis TB			Total	
		Ringan	Sedang	Berat		
IMT	Risiko Tinggi	Count	4	9	16	29
		Expected Count	7.5	9.3	12.2	29.0
		% within IMT	13.8%	31.0%	55.2%	100.0%
	Risiko Rendah	Count	9	7	5	21
		Expected Count	5.5	6.7	8.8	21.0
		% within IMT	42.9%	33.3%	23.8%	100.0%
Total	Count	13	16	21	50	

Expected Count	13.0	16.0	21.0	50.0
% within IMT	26.0%	32.0%	42.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	6.830 ^a	2	.033
Likelihood Ratio	6.998	2	.030
Linear-by-Linear Association	6.660	1	.010
N of Valid Cases	50		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 5.46.

Kondisi Imunokompromais * Tingkat Keparahan Spondilitis TB Crosstabulation

		Tingkat Keparahan Spondilitis TB			Total	
		Ringan	Sedang	Berat		
Kondisi Imunokompromais	Ya	Count	6	5	15	26
		Expected Count	6.8	8.3	10.9	26.0
		% within Kondisi Imunokompromais	23.1%	19.2%	57.7%	100.0%
	Tidak	Count	7	11	6	24
		Expected Count	6.2	7.7	10.1	24.0
		% within Kondisi Imunokompromais	29.2%	45.8%	25.0%	100.0%
Total	Count	13	16	21	50	
	Expected Count	13.0	16.0	21.0	50.0	
	% within Kondisi Imunokompromais	26.0%	32.0%	42.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	6.114 ^a	2	.047
Likelihood Ratio	6.288	2	.043
Linear-by-Linear Association	2.811	1	.094
N of Valid Cases	50		

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,24.

Riwayat TB Paru * Tingkat Keparahan Spondilitis TB Crosstabulation

		Tingkat Keparahan Spondilitis TB			Total	
		Ringan	Sedang	Berat		
Riwayat TB Paru	Ya	Count	6	11	12	29
		Expected Count	7.5	9.3	12.2	29.0
		% within Riwayat TB Paru	20.7%	37.9%	41.4%	100.0%
	Tidak	Count	7	5	9	21
		Expected Count	5.5	6.7	8.8	21.0
		% within Riwayat TB Paru	33.3%	23.8%	42.9%	100.0%
Total	Count	13	16	21	50	
	Expected Count	13.0	16.0	21.0	50.0	
	% within Riwayat TB Paru	26.0%	32.0%	42.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	1.514 ^a	2	.469
Likelihood Ratio	1.527	2	.466
Linear-by-Linear Association	.227	1	.633
N of Valid Cases	50		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 5.46.